



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

Three-Dimensional Spin Rotations at the Fermi Surface of a Strongly Spin-Orbit Coupled Surface System

Höpfner, P ; Schäfer, J ; Fleszar, A ; Dil, J H ; Slomski, B ; Meier, F ; Loho, C ; Blumenstein, C ;
Patthey, L ; Hanke, W ; Claessen, R

Abstract: The spin texture of the metallic two-dimensional electron system ($\sqrt{3}\times\sqrt{3}$)-Au/Ge(111) is revealed by fully three-dimensional spin-resolved photoemission, as well as by density functional calculations. The large hexagonal Fermi surface, generated by the Au atoms, shows a significant splitting due to spin-orbit interactions. The planar components of the spin exhibit a helical character, accompanied by a strong out-of-plane spin component with alternating signs along the six Fermi surface sections. Moreover, in-plane spin rotations toward a radial direction are observed close to the hexagon corners. Such a threefold-symmetric spin pattern is not described by the conventional Rashba model. Instead, it reveals an interplay with Dresselhaus-like spin-orbit effects as a result of the crystalline anisotropies.

DOI: <https://doi.org/10.1103/PhysRevLett.108.186801>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-62050>

Journal Article

Supplemental Material

Originally published at:

Höpfner, P; Schäfer, J; Fleszar, A; Dil, J H; Slomski, B; Meier, F; Loho, C; Blumenstein, C; Patthey, L; Hanke, W; Claessen, R (2012). Three-Dimensional Spin Rotations at the Fermi Surface of a Strongly Spin-Orbit Coupled Surface System. *Physical Review Letters*, 108:186801.

DOI: <https://doi.org/10.1103/PhysRevLett.108.186801>

Supplemental Material

Three-Dimensional Spin Rotations at the Fermi Surface of a Strongly Spin-Orbit Coupled Surface System

P. Höpfner, J. Schäfer, A. Fleszar, J. H. Dil, B. Slomski, F. Meier, C. Loho, C. Blumenstein,
L. Patthey, W. Hanke, and R. Claessen

This supplemental material provides details of the band-structure modeling according to Eq. (3):

$$H(k) = \left(\frac{\hbar^2 k^2}{2m^*} - C + c_h(k_+^6 + k_-^6) \right) \sigma_0 + v(k_x \sigma_y - k_y \sigma_x) + \lambda(k_+^3 + k_-^3) \sigma_z + i\zeta(k_+^5 \sigma_+ - k_-^5 \sigma_-),$$

where σ_0 is the 2×2 identity matrix, σ_x , σ_y and σ_z are Pauli matrices, $\sigma_{\pm} = \sigma_x \pm i\sigma_y$, and $k_{\pm} = k_x \pm ik_y$. This Hamiltonian is invariant with respect to symmetry operations of the C_{3v} group and time-reversal symmetry. For the following choice of parameters, the $E = 0$ energy surface of this model Hamiltonian reproduces well both, the Fermi contour and the spin structure of the metallic surface band in $(\sqrt{3} \times \sqrt{3})$ -Au/Ge(111): $m^* = 0.4 m_e$, $C = 0.77 \text{ eV}$, $c_h = 70 \text{ eV \AA}^{-6}$, $v = -0.1 \text{ eV \AA}^{-1}$, $\lambda = -7 \text{ eV \AA}^{-3}$, $\zeta = 5.5 \text{ eV \AA}^{-5}$.

The first part of the Hamiltonian, proportional to σ_0 , is the spinless part, giving rise to spin doubly degenerate states. The sixth-order in k term in the parentheses is responsible for the hexagonal warping of the constant energy contours. This term reproduces well the Fermi contour of our *ab-initio* calculations, before including the spin-orbit interaction.

The second part of the Hamiltonian, $v(k_x \sigma_y - k_y \sigma_x)$, is the well-known Rashba term. It has full $O(2)$ symmetry and produces an isotropic spin splitting. The two resulting constant-energy contours (corresponding to $S1_A$ and $S1_B$) have a spin texture being fully planar, vortical (clockwise for $S1_B$ and counter-clockwise for $S1_A$) and perpendicular to the surface momentum vector (k_x, k_y) .

The third part of the Hamiltonian, of third-order in k , was proposed by L. Fu in Ref. 24 in order to explain the hexagonal warping of the Dirac-fermion band at the surface of the topological insulator Bi_2Te_3 . Apart from affecting the shape of energy contours, it generates a z -component in the spin vector. This vertical part of the spin exhibits an alternating sign (*up* or *down*) between neighboring hexagonal sheets of the energy contours. The in-plane part of the spin vector remains, however, perpendicular to the momentum vector (k_x, k_y) . Along the $\bar{\Gamma}-\bar{K}$ azimuth, there is no z -component of spin due to symmetry reasons.

The last term, of fifth-order in k , $i\zeta(k_+^5 \sigma_+ - k_-^5 \sigma_-)$, was recently proposed by S. Basak *et al.* [25]. This term is needed to explain the *ab-initio* results showing radial rotations of the in-plane spin components for the Dirac-fermion bands at the surface of the topological insulator Bi_2Te_3 . Similarly as the third-order term, it contributes to the hexagonal warping of the energy contours, but its main function is the rotation of in-plane spin components towards a radial direction. This happens in close vicinity to the $\bar{\Gamma}-\bar{K}$ azimuth.




Figure S1: $E = 0$ energy contours derived from the model Hamiltonian with the in-plane spin texture of the spin-split states (a) $S1_A$ and (b) $S1_B$. The spin arrows are red for spins directed out-of-plane and blue for an orientation into the plane. Green color signifies fully in-plane spin alignment.

Figure S1 displays the energy contours and spin texture of the model Hamiltonian (Eq. (3)) for the choice of parameters listed above. For better clarity each spin-split state ($S1_A$ and $S1_B$) is shown in a separate plot.